




Assessment of Cerebrovascular Disease and White Matter Neurite Density in Alzheimer's Disease

Grant Roberts¹, Leonardo Rivera-Rivera², Kevin Johnson^{1,3}, Sterling Johnson², Douglas Dean III^{1,4}, Andrew Alexander^{1,5}, Oliver Wieben^{1,3}, and Laura Eisenmenger³

University of Wisconsin – Madison: Dept. of

¹Medical Physics, ²Medicine, ³Radiology, ⁴Pediatrics, ⁵Psychiatry

UNIVERSITY OF WISCONSIN-MADISON



Abstract #0267

15-20 May 2021

Declaration of Financial Interests or Relationships

Speaker Name: Grant S. Roberts

I have the following financial interest or relationship to disclose with regard to the subject matter of this presentation:

Company Name: GE Healthcare


Type of Relationship: Research Support




Background

- Alzheimer's disease (AD)
 - A β plaques and neurofibrillary tangles
 - Cortical atrophy
 - Typically thought of as disease of grey matter
- However, white matter (WM) alterations also occur¹⁻⁴
 - Likely vascular-mediated
 - Disrupts brain microcirculation
 - Impaired clearance of waste products
- We have also shown vascular changes present in AD⁵⁻⁷

1. Brun A, Englund, E (1986). *Ann Neurol* 19(3)
2. Agosta F, et al (2011). *Radiology* 258(3)
3. Slattery CF, et al (2017). *Neurobiol Aging* 57
4. Fu X, et al (2020). *Clin Neuroradiol* 30(3)
5. Rivera-Rivera LA, et al (2016). *JCBFM* 36(10)
6. Berman SE, et al (2015). *Neuroimaging* 1(4)
7. Rivera-Rivera LA, et al (2021). *JCBFM* 41(2)



Adapted From: Fu X, et al (2020). *Clin Neuroradiol* 30(3)



From: Rivera-Rivera LA, et al (2020). *JCBFM* 41(2)



Background

- Relationship between macrovascular flow and WM microstructure alterations is still unclear
- Goal:**
 - Utilize **4D Flow MRI** (cerebrovascular dynamics)
 - Utilize DTI **NODDI** (WM microstructure)
 - 20 AD and 41 cognitively normal (CN) subjects
- 1. Compare WM axon density between AD/CN groups
- 2. Correlate WM axon density with vascular measures
 - Carotid pulse wave velocity (stiffness)
 - Carotid pulsatility index (resistance)
 - Total cerebral blood flow

4D Flow




Image: Rivera-Rivera LA, et al (2016). *JCBFM* 36(10).




NODDI

ODI

NDI

f_{fw}



Methods

Patient Demographics

- 20 Alzheimer's disease subjects
 - Characterized as “dementia due to probable AD^{1,2}”
- 41 Cognitively normal subjects

	CN (N=41)	AD (N = 20)	p-value
Age (years)	74 ± 7	73 ± 9	0.96 ^a
Female (n, %)	23, 56.1	13, 65.0	0.58 ^b
Parental history of dementia (n, %)	1, 2.44	7, 35.0	0.001^b
APOE ε4 carrier (n, %)*	1, 2.44	6, 30.0	2.51e-04^b
SBP (mmHg)	132 ± 22	131 ± 19	0.78 ^a
DBP (mmHg)	78 ± 9	75 ± 6	0.23 ^a
HR (bpm)	62 ± 9	60 ± 11	0.55 ^a

CN = cognitively normal; AD = Alzheimer's disease; APOE = apolipoprotein E; SBP = systolic blood pressure; DBP = diastolic blood pressure; HR = heart rate

^aTwo sample t-test
^bFisher's exact test

Bold indicates statistical significance (p<0.05)

¹McKhann GM, et al (2011). *Alzheimers Dement* 7(3)

²Jack CR, et al (2018). *Alzheimers Dement* 14(4)





Methods


4D Flow Measurements

- Total cerebral blood flow (tCBF)
 - $tCBF = Flow_{ICA} + Flow_{BA}$
- Pulsatility index (PI)
 - Vascular resistance
- Pulse wave velocity (PWV)
 - Vascular stiffness
 - Local low rank reconstruction¹

MR Parameter	Value
Scanner	3.0T GE Discovery MR750
Coil	32 Channel Head
Sequence	PCVIPR ^{2,3}
Encoding Scheme	5-point balanced
Projections	11,000
TR	7.4 ms
TE	2.7 ms
V_{enc}	75 cm/s
Resolution	0.7 mm isotropic
Cardiac frames	100



UNIVERSITY OF WISCONSIN-MADISON



¹Rivera-Rivera LA, et al (2021). *JCBFM* 4(12).
²Gu T, et al (2005). *AJNR* 26(4).
³Johnson KM, et al (2008). *MRM* 60(6).




Methods

NODDI – Neurite Orientation Dispersion and Density Imaging¹

- Neurite density index (NDI)
- Orientation dispersion index (ODI)
- Free water fraction (f_{fw})

MR Parameter	Value
Scanner	3.0T GE Discovery MR750
Coil	32 Channel Head
Sequence	Spin-echo EPI
Shells	6 × $b=0$ s/mm ² 9 × $b=500$ s/mm ² 18 × $b=800$ s/mm ² 36 × $b=2000$ s/mm ²
Resolution	2 mm isotropic
TR	8575 ms
TE	76.8 ms
Flip angle	8 degrees




WISCONSIN UNIVERSITY OF WISCONSIN-MADISON

¹Zhang H, et al (2012). *Neuroimage* 61(4).

Methods

Diffusion Tensor Data

- FSL/MRtrix^{1,2}
- NODDI Matlab Toolbox³
 - NDI maps
- FSL
 - Tract-based Spatial Statistics: *tbss*⁴
 - Statistical Analysis: *Glm (randomise*⁵)



¹Jenkinson M, et al (2012). *Neuroimage* 62

²Tournier JD, et al (2019). *Nueroimage* 14

³Zhang H, et al (2012). *Neuroimage* 61(4)

⁴Smith SM, et al (2006). *Neuroimage* 31(4)


⁵Winkler AM, et al (2014). *Neuroimage* 92(100)



Methods

TBSS Hypothesis Tests

- Are there differences in NDI Between AD and CN subjects?
- Are there correlations between PWV/PI/tCBF and NDI for AD and CN subjects?
- *Post hoc* analysis in significant tracts
 - ROIs identified with JHU WM atlas
 - Mean NDI values were extracted for each subject



WISCONSIN
UNIVERSITY OF WISCONSIN-MADISON



Results




TBSS Hypothesis Tests

- Compare NDI Between AD and CN
 - CN NDI > AD NDI?

CN NDI > AD NDI

Similar results observed by Slattery et al¹ and Fu et al²



1. Slattery CF, et al (2017). *Neurobiol Aging* 57
2. Fu X, et al (2020). *Clin Neuroradiol* 30(3)



Results

ROI Analysis

- CN NDI > AD NDI
- WM regions identified on TBSS using JHU WM atlas
- ANCOVA
 - Adjusted for age/sex
 - Corrected for multiple comparisons

White Matter Regions Identified from TBSS (ICBM-DTI-81 Atlas Label #)	CN Mean NDI	AD Mean NDI	p-value
Genu of Corpus Callosum (3)	0.526 ± 0.041	0.507 ± 0.047	0.1329
Body of Corpus Callosum (4)	0.595 ± 0.028	0.573 ± 0.034	0.0137*
Splenium of Corpus Callosum (5)	0.635 ± 0.031	0.597 ± 0.032	5.2e-05**
Anterior corona radiata R (23)	0.473 ± 0.056	0.438 ± 0.054	0.0138*
Anterior corona radiata L (24)	0.474 ± 0.053	0.429 ± 0.054	0.0019**
Superior corona radiata R (25)	0.590 ± 0.046	0.556 ± 0.058	0.0113*
Superior corona radiata L (26)	0.583 ± 0.048	0.541 ± 0.059	0.0024*
Posterior corona radiata R (27)	0.512 ± 0.055	0.473 ± 0.063	0.0153*
Posterior corona radiata L (28)	0.505 ± 0.054	0.467 ± 0.049	0.0104*
Posterior thalamic radiation R (29)	0.515 ± 0.041	0.474 ± 0.041	3.9e-04**
Posterior thalamic radiation L (30)	0.487 ± 0.048	0.438 ± 0.047	3.3e-04**
Sagittal stratum R (31)	0.510 ± 0.030	0.492 ± 0.033	0.0558
Sagittal stratum L (32)	0.487 ± 0.035	0.467 ± 0.038	0.0541
External capsule R (33)	0.506 ± 0.030	0.485 ± 0.028	0.0075*
External capsule L (34)	0.509 ± 0.026	0.483 ± 0.028	2.9e-04**
Cingulate gyrus R (35)	0.536 ± 0.023	0.510 ± 0.022	7.8e-05**
Cingulate gyrus L (36)	0.542 ± 0.024	0.510 ± 0.023	8.4e-06**
Cingulum (hippocampus) R (37)	0.496 ± 0.029	0.463 ± 0.036	2.8e-04**
Cingulum (hippocampus) L (38)	0.494 ± 0.022	0.469 ± 0.025	3.5e-04**
Superior longitudinal fasciculus R (41)	0.592 ± 0.034	0.557 ± 0.056	0.0069*
Superior longitudinal fasciculus L (42)	0.594 ± 0.038	0.552 ± 0.053	7.8e-04**
Superior fronto-occipital fasciculus R (43)	0.557 ± 0.074	0.520 ± 0.061	0.0317*
Superior fronto-occipital fasciculus L (44)	0.542 ± 0.081	0.482 ± 0.070	0.0026*
Uncinate fasciculus R (45)	0.485 ± 0.026	0.454 ± 0.032	2.9e-05**
Uncinate fasciculus L (46)	0.484 ± 0.027	0.451 ± 0.022	1.6e-05**

Abbreviations: NDI=neurite density index; AD=Alzheimer's disease; CN=cognitively normal; TBSS=tract-based spatial statistics; R=right; L=left

*p-values <0.05; **p-value<0.002 (Bonferroni correction for 25 ROIs).



Results

TBSS Hypothesis Tests

- Compare NDI Between AD and CN
 - CN NDI > AD NDI?
- Correlation - PWV and NDI
 - No significant findings
- Correlation - PI and NDI
 - No significant findings

No significant findings!



WISCONSIN
UNIVERSITY OF WISCONSIN-MADISON




Results

TBSS Hypothesis Tests

- Compare NDI Between AD and CN
 - CN NDI > AD NDI?
- Correlation - PWV and NDI
 - No significant findings
- Correlation - PI and NDI
 - No significant findings
- Correlation - tCBF and NDI
 - CN $\beta_{tCBF} \neq 0$?
 - ~~AD $\beta_{tCBF} \neq 0$?~~

CN $\beta_{tCBF} > 0$



Results

ROI Analysis

CN $\beta_{tCBF} > 0$

- Multiple Regression
 - Adjusted for age/sex
 - Corrected for multiple comparisons





White Matter Regions Identified from TBSS (ICBM-DTI-81 Atlas Label #)	CN $\beta_{tCBF} > 0$		
	β_{tCBF}	R ²	p-value
Genu of corpus callosum (3)	0.025	0.129	0.019*
Body of corpus callosum (4)	0.025	0.247	4e-04**
Splenium of corpus callosum (5)	0.019	0.176	0.013*
Anterior corona radiata R (23)	0.026	0.283	0.045*
Anterior corona radiata L (24)	0.027	0.212	0.034*
Superior corona radiata R (25)	0.019	0.237	0.075
Superior corona radiata L (26)	0.015	0.174	0.203
Posterior corona radiata R (27)	0.036	0.245	0.007*
Posterior corona radiata L (28)	0.035	0.216	0.008*
Posterior thalamic radiation R (29)	0.020	0.206	0.049*
Posterior thalamic radiation L (30)	0.021	0.169	0.078
Superior long. fasciculus R (41)	0.028	0.349	0.001**
Superior long. fasciculus L (42)	0.023	0.221	0.012*

Abbreviations: tCBF = total cerebral blood flow; NDI=neurite density index; CN=cognitively normal; AD=Alzheimer's disease; TBSS=tract-based spatial statistics
 β : regression coefficient; R²: adjusted coefficient of determination

*p-values <0.05; **p-value<0.004 (Bonferroni correction for 13 ROIs).



Results



Discussion

1. Slattery CF, et al (2017). *Neurobiol Aging* 57
2. Fu X, et al (2020). *Clin Neuroradiol* 30(3)

Significant Findings

1. White matter density was decreased in AD subjects
 - Variety of WM regions^{1,2}
2. Positive correlation between cerebral blood flow and axon density in CN subjects
 - Corpus callosum
 - Superior longitudinal fasciculus
3. No associations between CVD and NDI for AD subjects

Limitations and Future Work

- Small sample size
- Need longitudinal data with subjects along the AD continuum to look at potential *causative* affects of CVD on WM microstructure

Conclusion

- WM microstructure alterations, as measured by NDI, were observed in AD group
- Cerebral blood flow was significantly correlated with WM axon density only in control group



Acknowledgements



- Wieben Lab
- Eisenmenger Lab
- ADRC Vascular Team
- Waisman Brain Imaging Team
- GE Healthcare

Funding: F31AG071183; KL2TR002374; UL1TR002373;
AARFD-20-678095; P50AG033514; RO1AG021155

